# A Key to the Bats of the Philippine Islands

Nina R. Ingle and Lawrence R. Heaney

#### Abstract

An identification guide is presented for the six families and 70 species of bats now known from the Philippine Islands, based on a key and a set of standardized measurements. Most critical characters are illustrated, and detailed drawings are provided of the skulls of 42 species.

#### Introduction

The Philippine Islands (fig. 1) support a large and diverse fauna of mammals: over 170 species are now known, compared, for example, to the 105 species known from Madagascar, which has nearly twice the land area (Heaney et al., 1987; Jenkins, 1987). About 100 species are endemic to the Philippines, giving the country an unusually high number and percentage of unique species (Hauge et al., 1986; Heaney, 1986, 1991; Heaney et al., 1987; Koopman, 1989).


One of the most diverse and, in general, poorly known mammalian orders in the Philippines is the Chiroptera. Sixty-eight species of bats were known from the Philippines when the last checklist was prepared (Heaney et al., 1987), and two more are now known (*Pteropus dasymallus* and *Harpiocephalus harpia*). In number of species, bats exceed even rodents, of which 67 are now known (Heaney et al., 1987; Musser and Heaney, 1992). By our best estimate, 22 species of bats, about 31%, are endemic to the Philippines, again an unusually high number (Heaney, 1991; Koopman, 1989).

The high levels of species richness and endemism are factors of special importance currently because of the rapid rate of loss of natural habitat in the Philippines. Roughly 94% of the Philippine land area was once covered by forest; that figure had been reduced to 40% at the end of World War II, and current estimates of forest cover range from 25% to less than 20%, depending in part on the amount of degraded forest that is included (Collins, 1990; Hauge et al., 1986; Myers, 1988; Utzurrum, 1991). The ongoing forest destruction

poses an especially grave problem because many species of bats, especially endemics, depend primarily on forest (Heaney et al., 1987; Heaney and Utzurrum, 1991; Heideman and Heaney, 1989). Two species of bats (*Acerodon lucifer* and *Dobsonia chapmani*) are believed to have become extinct in the last 100 years, and many others are threatened (Heaney and Heideman, 1987; Heaney et al., 1987).

No identification guide to the 70 species of bats recorded from the Philippines currently exists. The most recent work describing Philippine bats, Taylor's "Philippine Land Mammals", was published in 1934 and does not include the many species subsequently described or recorded from the Philippines, nor does it reflect the many changes in taxonomy that have taken place over nearly 60 years. Moreover, Taylor's keys were intended for use with museum study specimens, not live animals in the field. Thus, identification of Philippine bats in recent years has been primarily by comparison with museum specimens. This requires access to a comprehensive reference collection and is not an option for researchers in many places.

This key is intended to permit identification of all bat species that have been recorded from the Philippines, to the extent that current knowledge permits. It is our hope that this key will encourage more research on the Philippine bat fauna, which, with its diversity and distribution over many islands with differing habitats and climates, serves as an excellent subject for studies of biogeography and many aspects of ecology. We also hope that a greater knowledge about Philippine bats will contribute to efforts toward their conservation.






Fig. 2. Generalized pteropodid bat, with important structures indicated.

### Methods

The descriptions and measurements in this key have been based primarily on examination of specimens in the American Museum of Natural History (AMNH), British Museum (Natural History) (BMNH), Delaware Museum of Natural History (DMNH), Field Museum of Natural History (FMNH), Royal Ontario Museum (ROM), Philippine National Museum (PNM), Silliman University Museum of Natural History (su), University of Michigan, Museum of Zoology (UMMZ), and United States National Museum of Natural History (USNM). Our descriptions and measurements are based on Philippine specimens except in a few cases involving species that are rare in collections; in those cases, specimens from elsewhere or characters from published accounts were utilized.

For standard external measurements we relied preferentially on data from animals that we had collected, but we also utilized data on specimen labels and measured specimens in collections. We took cranial measurements with dial or digital calipers following deBlase and Martin (1974) and Heaney and Peterson (1984); these measurements are described briefly in the following section, Anatomy and Measurements of a Bat.

We consulted a variety of publications as aids in building the keys and descriptive sections, relying heavily on Harrison (1966), Hill (1983), Lawrence (1939), Lekagul and McNeely (1977), Medway (1969), Miller (1907), Payne et al. (1985), and Taylor (1934). These publications, the references cited in Hill (1983), and publications referred to in the sections for each bat family should be consulted for descriptions of characters not treated in the key.

# Anatomy and Measurements of a Bat

Familiarity with basic terminology on the anatomy of a bat is necessary to use this key; Figure 2 shows the most important external structures. The external ear (pinna) has two structures that can be useful for identification. The tragus is a

Fig. 1. Map of the Philippines. Recent islands are outlined by continuous lines, and the extent of late Pleistocene islands is indicated by the shaded areas.




Fig. 3. Left external ears (pinnae) of representative bats. A, Pteropodidae; B, Rhinolophidae; C, Vespertilionidae.

projection from the base of the ear (fig. 3C); this structure is absent in members of the Pteropodidae (fig. 3A) and Rhinolophidae (fig. 3B) and is quite small in the Molossidae. The antitragus is the flap along the lower posterior margin of the ear (figs. 3B, 12); it is especially well developed in rhinolophids and most molossids. Rhinolophids and megadermatids have noseleaves, structures formed from elaborate folds of skin on their noses (figs. 8B,C, 12); bats of other families have no such structure.

All bats have four types of teeth: incisors, canines, premolars, and molars (fig. 5). All bats possess a pair of large, sharp, conical teeth called canines on the upper and lower jaws. The smaller teeth at the front of the mouth, anterior to the

canines, are incisors. Posterior to the canines are the premolars and molars, collectively called cheek teeth. The large teeth in this series are referred to as molariform teeth. Molariform teeth are usually low and broad with cutting or grinding surfaces; the anterior cheek teeth are tiny pegs in some species.

External and cranial measurements provide substantial aid in identification; we have therefore included tables containing typical adult measurements. The definitions of standard external measurements are illustrated in Figure 4. The cranial measurements included in the tables are shown in Figure 5 and are defined as follows: condylobasal length (CBL), distance from the posterior edge of the occipital condyles to the anterior tip of the premaxillaries; condylocanine length (CCL), distance from the posterior edge of the occipital condyles to the anterior edge of the base (alveolus) of the canines; and maxillary toothrow (C1 to last M), distance from the posterior edge of the last upper molar to the anterior edge of the upper canine, taken along the bone line of the alveolus.

### Use of the Key

Using the key most effectively requires several steps that are explained in detail in the following sections. First, the bat should be identified as to




Fig. 4. Bat with external measurements indicated.




Fig. 5. Bat skull with teeth and cranial measurements indicated.

age and sex. It should then be identified to family with the Key to Families, and then to species with the key to that family. Next, all external measurements should be taken and matched against those in the tables of measurements; if the measurements do not match, the identification is questionable and the process should be begun again from the first step. Finally, if a skull is available, it should be visually compared with Figures 19–60 to see if it matches the features of that species or genus.

### Determination of Age and Sex

The age of bats can be estimated by the degree of ossification of the joints in the digits of the wing (Anthony, 1988). In juvenile bats, the joints have cartilaginous discs where growth takes place. When a light is shown through the wing of a live bat, the bands of cartilage at the joints appear partly translucent (fig. 6B). In adult bats, the bones are fully ossified and the joints appear opaque (fig. 6A). The shape of the joints of the digits also differs between juvenile and adult bats; this is especially useful in determining the age of animals prepared as stuffed skins or fluid-preserved specimens. In juvenile bats

the joints are swollen and tapered (fig. 6B), whereas in adult bats the joints are knobby and more distinct from the bone shaft (fig. 6A).

Bats can be sexed by examination of the external genitalia. Males have a conspicuous penis (except




Fig. 6. Joints in digits of wing. Stippled areas represent bone and open areas represent cartilage. A, adult; B, juvenile.

subadult male *Rousettus*, in which the penis is retracted into the abdomen). Both sexes possess axillary nipples on the upper chest, usually near the armpit (axilla). However, the nipples of adult females are more prominent than those of males. In female rhinolophids and megadermatids, a pair of inguinal papillae that look like nipples is present just anterior to the genitals.

# **Identification to Family and Species**

Following determination of age and sex, the bat should be identified to family with the Key to Families, and then to species with the key for the appropriate family. The keys consist of sets of alternatives arranged in order of increasing restrictiveness. Alternatives belonging to the same set are preceded by the same number with different superscripts (e.g., 1, 1', 1"). For each set, the user should choose the best alternative; this indicates either the next set of alternatives to be considered. or the name of the family or species. All distinguishing characters should be checked carefully before a choice is made. This will minimize the likelihood of incorrectly identifying a bat. Researchers should bear in mind that this key includes only those species now known from the Philippines, and that discovery of undescribed species and previously unreported species is likely.

Figures illustrating important descriptive characteristics are referred to in the key and should be consulted. Fur coloration can be useful in identification; it should be observed on a dry animal or study skin.

#### Use of Measurement Tables

Once a bat has been identified to species, it is good practice to check if its measurements agree with those given in the tables. Cranial measurements should be taken with calipers. External measurements can be taken with a ruler, although calipers provide more accurate measurements of forearm length and of small structures such as parts of the noseleaves of rhinolophids.

Because forearm length is a very useful measurement for identifying adult bats, ranges for forearm length are given in the keys for each species, usually at the end of the final description for that species. Note that forearm lengths are given even

when overlap in the ranges of two or more species means that the measurement cannot be used to fully distinguish between the species. When other measurements are particularly useful for separating similar species, they are also given in the key. Measurements should not be used to identify juvenile bats, although hind foot length is sometimes useful because the feet of juvenile bats approach adult size long before most other structures have ended growth.

### Comparison with Skull Drawings

Skulls are very important in identification of bats. Thus, although we have written this key with the intention that live animals and preserved specimens can be identified without examination of their skulls, we have included skull drawings of common Philippine bats (figs. 19-60). The drawings will aid in identification of museum specimens with cleaned skulls and will provide a means of verifying an identification based on external characteristics. The skull drawings also serve as a guide to head shape. Even if use of the key yields an apparently unambiguous identification, it is always best if a skull can be examined from a voucher specimen. The skull should be compared with the figures, and cranial measurements should be taken and compared with those in the tables. Dentition, although most easily visible on a skull, can also be observed, at least partially, on a live animal and should be examined when possible. The mouth can be gently pried and held open by a toothpick, and the lips can be lifted to view the cheek teeth. With small bats a magnifying lens is a great help. Note that if the jaws are closed, some teeth, particularly the canines, may obstruct the view of others.

Users of this key should not overlook the fact that variation in both quantitative and qualitative characters is present in all species. Whenever possible, samples of more than one individual for each species should be examined and the most common character states used in the key.

The identities of a few species of Philippine bats that are extremely poorly represented in collections (sometimes only by the holotype) are uncertain. When possible, these species have been included in the key on the basis of whatever descriptions are available, and further comments on their identification are provided in the Notes section. We also comment briefly on two new records of bats for the Philippines in this section.

# Collecting and Preserving Voucher Specimens

In any study, several individuals of each species should be preserved as voucher specimens and deposited in a museum collection. Voucher specimens are necessary to verify identification, since many species can be identified definitively only after careful study of the external and cranial morphology. Such specimens are invaluable for adding to our current, very limited knowledge of Philippine bats, particularly of distribution and interand intraspecific variation. Specimens should be collected and prepared carefully, following all regulations for the scientific study of wildlife.

Bats are most easily preserved in fluid. They should be killed in a quick and painless manner, examined for reproductive condition, and weighed, and their standard external measurements taken (total length, tail length, ear length, and hind foot length). The date, habitat, collection locality, elevation, collector, collector's field number, and

external measurements should be recorded in India ink (which is best) or pencil (which is usually acceptable), not with other kinds of ink (which are not permanent). These data should accompany the specimen as part of the permanent record. The specimen should have a permanent label attached that bears the collector's name and field number; other data may go onto the same label or into a field catalog. The specimens should then be rinsed lightly with soapy water, injected with a 10% solution of formalin, and immersed in a 10% solution of formalin. After about three days, they can be rinsed with clean water and transferred to 70% alcohol (methanol is best) for permanent storage. Skulls can be extracted and cleaned in the museum.

The Philippine National Museum, University of the Philippines at Los Baños, Silliman University, and a number of other institutions maintain research collections and accept voucher specimens for their permanent collections, and the staff may be able to help with identifications.

## **Key to Families of Philippine Bats**

- 1. Interfemoral membrane is absent or reduced, forming narrow margin along insides of legs; tail short (up to 20% of body length) or absent, never completely enclosed by interfemoral membrane (fig. 7A); both thumb and second finger with claw (except *Eonycteris* and *Dobsonia*; fig. 2); both tragus and antitragus absent (figs. 3A, 8A); ear margin forms continuous ring ... Megachiroptera: Pteropodidae
- l'. Interfemoral membrane is a continuous expanse of skin stretching between legs (figs. 7B-F; except Coelops, which has a concave posterior margin to the membrane and no tail); tail present (except Megaderma and Coelops, which both possess noseleaves, structures not present in any pteropodids), usually comprising more than 20% of body length; the second finger does not have a claw; either tragus (fig. 3C) or antitragus (fig. 3B) or both present; ear margin does not form continuous ring ...

- 4". Tail projects beyond posterior margin of interfemoral membrane for over half its length (fig. 7E); ears thick and fleshy (fig. 8E); anterior edge of bony palate continuous, not emarginate ..........




Fig. 7. Tails of representatives of Philippine bat families. A, Pteropodidae (Rousettus); B, Megadermatidae (Megaderma); C, Vespertilionidae (Miniopterus); D, Rhinolophidae (Rhinolophus); E, Molossidae (Chaerephon); F, Emballonuridae (Taphozous).


# Family Pteropodidae: Fruit Bats

All fruit bats are characterized by dog-like heads (fig. 8A), with eyes that are proportionately larger

than those of most other bats; the large eyes are associated with their dependence on vision for orientation and their lack of echolocation (sonar) systems (except for the very simple and limited one used by *Rousettus*). They do not have the elaborate

FIELDIANA: ZOOLOGY

Fig. 8. Heads of representatives of Philippine bat families (not to same scale). A. Pteropodidae (*Rousettus*). Note that the ear margin is continuous, neither tragus nor noseleaf is present, the eyes are large, and the face is dog-like. B. Megadermatidae (*Megaderma*). Note that a noseleaf is present, the tragus is forked, and the large ears are connected across the top of the forehead. C. Rhinolophidae (*Rhinolophus*). Note the elaborate noseleaf, the well-developed



antitragus, and the absence of a tragus. D. Emballonuridae (*Taphozous*). Note that there is no noseleaf and that a tragus is present. E. Molossidae (*Chaerephon*). Note that there is no noseleaf and that the ears are thick and fleshy. F. Vespertilionidae (*Miniopterus*). Note that there is no noseleaf and that a tragus is present.

folds of skin in the nasal region (also associated with the use of echolocation) that are present in two other families of bats in the Philippines (Megadermatidae and Rhinolophidae). The external portion of the ear is simple and of moderate size, and the margin of the ear forms a continuous ring. Fruit bats do not have a tragus or antitragus (figs. 3A, 8A). All but three species (Dobsonia chapmani, Eonycteris robusta, and E. spelaea) have a claw on the second digit (fig. 2), along the leading edge of the wing; bats belonging to other families lack this claw. Many species have no tail, and in none does the tail comprise more than 20% of the body length. Their size varies greatly, with forearm lengths ranging from 40 to 215 mm, and weight from about 16 g (Alionycteris paucidentata) to at least 1,200 g (Acerodon jubatus; table 1).

Most species show substantial sexual dimorphism; males are larger than females and often have broader, more heavily muscled heads. Males of many species possess a ruff of fur (often colored yellow or rusty red) around the neck and on the shoulders that is lacking or poorly developed in females. Subadults of both sexes resemble adult females. Parous adult females of all species have a pair of axillary nipples that are large and prominent; males and subadult females have very small but visible nipples.

The skulls of all pteropodids are characterized by the presence of prominent postorbital processes, and all but a few species have postorbital foramina (figs. 19–32). The dental formula is variable. Most species have broad, blunt, rounded molariform teeth, but the nectarivorous species have greatly reduced premolars and molars. All species have long, prominent canines. In many species, males have longer canines than do females and often have more strongly developed cranial crests.

As the English name for the family implies, most fruit bats feed on fruit, but three species (Eonycteris robusta, E. spelaea, and Macroglossus minimus) feed primarily on nectar and pollen. Fruit bats often are abundant, being uncommon only in upper montane and mossy forest (Heaney and Rickart, 1990; Heaney et al., 1989). In forested areas, small to medium-sized species roost in hollow trees and in foliage, either alone or in groups; the large flying foxes roost in exposed treetops. Several species roost in caves, often forming colonies in the hundreds or thousands.

Andersen (1912) provided descriptions of most species of pteropodids found in the Philippines. Additional descriptions may be found in Bergmans (1975, 1978), Francis (1989), Heaney and Peterson (1984), Klingener and Creighton (1984), Kock (1969a,b,c), Musser et al. (1982), Peterson and Fenton (1970), Rookmaaker and Bergmans (1981), and Yoshiyuki (1979).

### Key to Pteropodidae

1. Claw on thumb but not on second digit
1'. Claws on both thumb and second digit (fig. 2)
2. Wings attach to body along midline of back; two upper incisors and two minute lower incisors;
forearm 123-131 mm; skull as in Figure 21
2'. Wings attach along sides of body; four upper and four lower incisors, all very small; forearm 67-
82 mm( <i>Eonycteris</i> ) 3
3. Pair of prominent 2-6-mm-long kidney-shaped glands lateral to anus; tail 12-20 mm; forearm 67-
80 mm; skull as in Figure 23 Eonycteris spelaea
3'. No glands near anus; tail 20-28 mm; forearm 67-82 mm; skull as in Figure 22
Eonycteris robusta
4. Forearm 94–215 mm; tail absent; four upper and four lower incisors
4'. Forearm 41-92 mm; tail present or absent; number of incisors varies between species 13
5. Wings with prominent pale blotches, particularly along thumb and anterior edge of wing adjacent
to first and second digit, especially at wing tips; forearm 135-141 mm Pteropus leucopterus
5'. Wings dark brown without prominent pale blotches; forearm 94-215 mm
6. Pelage on dorsal surface of lower back pale brown or gray
6'. Pelage on dorsal surface of lower back dark brown or black, sometimes with yellow flecks 8
7. Forearm 132-165 mm; condylobasal length 65-70 mm; three cusps on second and third upper
molariform teeth, including a well-developed anterolingual cusp
7'. Forearm 94-113 mm; condylobasal length 46-52 mm; two cusps on second and third upper mo-
lariform teeth (no anterolingual cusps); skull as in Figure 30 Pteropus pumilus

TABLE 1. Measurement ranges of adult Philippine Pteropodidae. Ranges represent at least 10 individuals, except when sample sizes are given in parentheses. Measurements, as defined in text, were taken from Philippine specimens.

Species	CBL	C'-last M	Total Length	Tail	Ear	Forearm	Weight
Acerodon jubatus	72.5–85.5	29.0–36.0	255-310	0	31-41	165-215	900-1.250
Acerodon leucotis	65.0-70.0	25.5-28.0	220-250	0	29–32	132-165	1
Acerodon lucifer	71.0–73.5	29.0–31.0	ı	0	1	167-170	ı
Alionycteris paucidentata	21.0-22.0	6.8-7.3	(5) 64–70	(5) 0	(5) 11–12	(5) 45–50	14-18
Cynopterus brachyotis	26.0-29.5	8.3-10.0	88-110	3-15	14-20	58–68	26-41
Dobsonia chapmani	48.5-53.5	19.0-21.0	218-221	23-26	25-27	123-131	1
Dyacopterus spadiceus	(2) 41.3–42.2	(2) 14.4 - 15.3	(1) 171	(1) 29	(1) 22	(1) 92	ı
Eonycteris robusta	33.9–37.9	11.6–14.0	127-155	20-28	19-23	67–82	75-80
Eonycteris spelaea	31.3–35.6	10.8-13.1	121-145	12-20	18-23	67–80	48-90
Haplonycteris fischeri	22.0-25.4	7.6-9.2	65-80	0	11-16	46-53	16-21
Harpyionycteris whiteheadi	39.7-43.3	15.4–16.6	130-156	0	19–25	80-91	99-122
Macroglossus minimus	24.7–26.5	9.6-9.2	65-81	0	12-18	4145	15-20
Megaerops wetmorei	21.7–22.9	7.1–7.7	20-16	9	12-15	45-52	17-20
Nyctimene rabori	33.4–35.3	11.2-12.1	131-149	20–30	17-24	71–79	61–74
Otopteropus cartilagonodus	20.8-23.0	7.0-8.0	70-76	0	12-14	43–50	15-17
Prenochirus jagori	32.5–37.2	11.0-13.3	120-145	6-18	18-25	06-92	62-97
Ptenochirus minor	27.2–31.7	9.5-11.0	98-125	4-12	13-22	82-09	31–62
Preropus dasymallus	57.6-64.2	22.3–24.6	220-230	0	(1) 25	(3) 133–152	(3) 380-490
Preropus hypomelanus	56.0-65.5	21.4–24.6	210-240	0	28-32	136-149	425-450
Pteropus leucopterus	56.5-63.0	16.5-22.0	205-240	0	26-28	135-141	340
Preropus pumilus	46.0-51.5	16.5-19.5	155-180	0	20-28	94-113	145-200
Preropus speciosus	53.9-61.5	20.3–22.9	190-215	0	24-30	118-133	ı
Pieropus vampyrus	68.8-79.0	26.0-31.5	270-320	0	35-45	179-204	725-810
Rousettus amplexicaudatus	34.0-38.8	12.1–14.4	128-154	13-24	18-24	80-92	64-106




Fig. 9. Frontal views of fruit bat skulls, showing incisors and canines (not to same scale). A, Otopteropus cartilagonodus (Alionycteris paucidentata and Haplonycteris fischeri have an equal number of incisors); B, Cynopterus brachyotis; C, Nyctimene rabori; D, Ptenochirus minor (Ptenochirus jagori incisors are very similar).

8.	Forearm 165–215 mm; condylobasal length 68–86 mm; fur on upper back sometimes completely black
8′.	Forearm 118–152 mm; condylobasal length 54–66 mm; fur on upper back never completely black, usually golden
	Dorsal pelage may be completely dark brown or black; if upper back is golden, the posterior margin of the golden area forms a sharply defined transverse line with the dark brown lower back; tips of ears nearly pointed; two cusps on second and third upper molariform teeth (no anterolingual cusps); condylobasal length 68–79 mm; forearm 179–204 mm; skull as in Figure 31 Pteropus vampyrus
	Dorsal pelage never completely dark brown or black; a golden patch on top of head, extending anterior to line between ears, is always present; the golden dorsal pelage never forms a transverse line along the edge of the dark brown of the lower back; tips of ears bluntly rounded; three cusps on second and third upper molariform teeth, including a well-developed anterolingual cusp; forearm 165–215 mm; condylobasal length 71–86 mm
10.	Forearm 167–170 mm; condylobasal length 71–74 mm (occurs only on Panay Island; see Notes)
10'.	Forearm 165–215 mm; condylobasal length 72–86 mm; skull as in Figure 19 Acerodon jubatus
11.	Forearm 118–133 mm; condylobasal length 53–62 mm (occurs only from Zamboanga to Sulu)
11'.	Forearm 133–152 mm; condylobasal length 56–66 mm
12.	Lower legs heavily furred nearly to ankle; forearm 133–152 mm (see Notes)
	Lower legs nearly naked; forearm 136–149 mm; skull as in Figure 29 Pteropus hypomelanus
13.	Tail absent
	Tail present (fig. 7A; it may be small, so look carefully)
14.	50 mm
14'.	No lumps of soft white tissue at margins of ears; forearm 41–91 mm
	Forearm 80–91 mm; well-developed secondary cusps on canines (one on upper canines, two on lower canines); six cusps on last two upper and lower molariform teeth; dorsal surface of hind feet thickly furred; skull as in Figure 25
15.	Forearm 41–53 mm; canines do not have well-developed secondary cusps; two or three cusps on molariform teeth; dorsal surface of hind feet may or may not be thickly furred
16.	Muzzle long and slender (fig. 10A); nostrils are not tubular and do not project beyond rest of muzzle; wing membrane attaches on top of foot, above gap between third and fourth toes from outside (fig.




Fig. 10. Dorsal views of fruit bat muzzles (not to same scale). A, Macroglossus; B, Haplonycteris; C, Nyctimene; D, Rousettus; E, Cynopterus.

11A); teeth, except canines, greatly reduced; forearm 41-45 mm; skull as in Figure 26 ......

16'.	Muzzle short and broad; nostrils tubular (as in fig. 10B); wing membrane attaches on side of foot
	(on or above outermost toe; fig. 11C); teeth robust; forearm 43–53 mm
17.	Four upper and two lower incisors (fig. 9D); forearm 45-52 mm Megaerops wetmorei
17'.	Two upper and two lower incisors (as in fig. 9A); forearm 43–53 mm
18.	Band of pale fur along dorsal surface of forearm; interfemoral membrane present; forearm 46–53
	mm; skull as in Figure 24
18'.	No band of pale fur along dorsal surface of forearm; no interfemoral membrane; forearm 45-50
	mm
19.	Ears and skin on dorsal surface of bones of wings with prominent pale yellow spots; dark stripe
	along most of dorsal midline; two upper and no lower incisors (fig. 9C); nostrils elongated into tubes
	about 2-3 mm long (fig. 10C); forearm 71-79 mm; skull as in Figure 27 Nyctimene rabori
19'.	Ears and wings without yellow spots; no stripe along dorsal midline; four upper incisors and at
	least two lower incisors; nostrils not elongated into tubes (although when viewed from above they
20	may appear slightly tubular); forearm 45–92 mm
20.	Muzzle moderately long and tapered (fig. 10D); anterior surface of upper canines with vertical
	groove (not always prominent); wing membrane terminates above gap between outermost toe and
	second toe from outside (fig. 11B); forearm 80–92 mm; skull as in Figure 32
201	Muzzle short and broad (fig. 10E); anterior surface of upper canines smooth, not grooved; wing
20.	membrane terminates either on outermost toe (fig. 11C) or above gap between outermost toe and
	second toe from outside (fig. 11B); forearm 45–92 mm
21.	Four upper and two lower incisors (outer pair of upper incisors much smaller than inner pair; fig.
	9D)
21'.	Four upper and four lower incisors (fig. 9B)
	Forearm 76–90 mm; skull as in Figure 28

# Family Emballonuridae: Sheath-Tailed Bats

Sheath-tailed bats are small to moderate-sized bats (forearm 44–71 mm; table 2). They are the only bats in the Philippines in which the tail perforates the dorsal surface of the interfemoral membrane (fig. 7F), so that the basal portion (usually about half) of the tail is enclosed within the membrane and the distal portion is free, lying on the dorsal surface of the membrane. The eyes are small to moderate in size, and the ears are simple and of moderate size (fig. 8D). A tragus is always present. The muzzle is rather pointed, with the nostrils at the tip; it does not have a noseleaf.

The skulls of emballonurids have postorbital processes that are very well developed (figs. 33–35); these processes are absent in vespertilionids, which are superficially the most similar to emballonurids. Emballonurid premaxillaries are small and delicate; they do not fuse to each other at the anterior midline and are attached to the maxillaries by flexible connective tissue. The anterior edge of the palate has a deep emargination.

All of the sheath-tailed bats in the Philippines feed on insects. Common roosting sites are caves, hollow trees, and attics of buildings; they usually aggregate in moderate to large groups (ten to several hundred).

### Key to Emballonuridae

# Family Megadermatidae: False Vampire Bats

This is the least diverse of bat families in the Philippines, with only a single species present, *Megaderma spasma* (fig. 8B). *M. spasma* is a medium-sized bat (forearm 57–63 mm; table 2) easily recognized by its large ears that meet across the top of the forehead, erect, simple noseleaf, and lack of tail (although the interfemoral membrane is well developed).

The skull of *Megaderma spasma* is of moderate size and its stoutly constructed (fig. 36). The premaxillaries are absent, so there are no upper in-

cisors, and the anterior edge of the palate has a deep emargination. The postorbital processes are very poorly developed. The upper canines have a large secondary cusp.




FIG. 11. Posterior attachment of wing membrane on left foot. A, *Macroglossus*; B, *Rousettus*; C, *Ptenochirus/Haplonycteris*.

Table 2. Measurement ranges of at least 10 individuals of adult Philippine Emballonuridae and Megadermatidae. Measurements, as defined in text, were taken from Philippine specimens.

Species	CBL	CCL	C¹-last M	Total Length	Tail	Hind Foot	Ear	Fore- arm	Weight
Emballonura alecto	13.3-14.2	12.5-13.4	5.0-5.8	56-69	9-12	7-10	12-16	44-49	4-6.5
Saccolaimus saccolaimus	20.9-23.3	_	8.8-9.9	103-117	20-28	15-19	16-20	66-71	28-36
Taphozous melanopogon	19.6-20.6	_	8.1-8.8	100-111	20-25	11-15	20-24	61-65	20-29
Megaderma spasma	_	21.2-23.6	8.7–9.4	70–89	0	18-22	36-43	57–63	21–27

Megaderma spasma feeds primarily on large insects (especially cicadas and katydids) and occasionally on small vertebrates such as lizards, frogs, and small birds (D. Balete, pers. comm.). It has been found roosting in caves and in hollow trees, singly and in small groups.

Because only a single species is present, no key to the family is necessary.

### Family Rhinolophidae: Horseshoe and Leaf-Nosed Bats

The family Rhinolophidae is composed of about ten genera and over 100 species; it is represented in the Philippines by three genera (Coelops, Hipposideros, and Rhinolophus) and at least 17 species (see Notes). All rhinolophids have a noseleaf, a structure consisting of elaborate folds of skin in the nasal region. The shape of the noseleaf varies among species and is an important character for identification (fig. 12). In members of the genus Rhinolophus, the posterior noseleaf is long and pointed (fig. 12A) and the anterior noseleaf is horseshoe-shaped. Between the anterior and posterior noseleaves lies the sella, an anterior-facing structure that is connected to the posterior noseleaf by the connecting process. In some species, supplementary leaflets are present lateral and ventral to the posterior noseleaf (fig. 12A). In Coelops and Hipposideros, the posterior noseleaf is low and rounded (fig. 12B) and may be divided into pockets by vertical septa. The intermediate noseleaf is a cushion-like structure. Supplementary leaflets are sometimes present lateral and ventral to the anterior noseleaf (not shown in fig. 12B, but see fig. 12A). Between the nostrils is the internarial septum. Rhinolophids lack a tragus, but many species, especially those belonging to the genus Rhinolophus, possess a well-developed antitragus, a flap of skin on the lower posterior margin of the ear (fig. 12). Most rhinolophids are small (forearm 34-60 mm), but a few are medium-sized (forearm up to 90 mm; table 3).


The skulls of most rhinolophids are delicate and slender, although some, especially those of the larger Hipposideros, are fairly stout. Postorbital processes are absent. Premaxillary bones are present but are small and connected to the skull by cartilaginous articulations, and so are movable in live or freshly killed animals. Incisors are present but are moderate to very small. The lower incisors are trilobed, although this is often inconspicuous. Many species have a strongly expanded nasal region (which supports the noseleaf structures), giving the skull a strongly sinuous dorsal profile in lateral view (figs. 37-46).

Rhinolophids are insectivorous. Some species roost in caves, in numbers ranging from a few to several hundred. Other species roost in hollow trees, hollow fallen logs, and other sites in forest. Although rhinolophids usually comprise a small proportion of captures in mist nets, at times a single species may be the most common bat species netted at a given site.

For further descriptions of members of the genus Hipposideros, see Hill (1963a) and Jenkins and Hill (1981). For Rhinolophus, see Andersen (1905a,b,c,d).

#### Key to Rhinolophidae

1. Posterior noseleaf pointed (fig. 12A); six pairs of lower cheek teeth ............ (Rhinolophus) 2 3. Dorsal tip of connecting process sharply pointed (fig. 13A); forearm 47–49 mm .....






Fig. 12. Noseleaves of Rhinolophus and Hipposideros. A, Rhinolophus; B, Hipposideros.

3'.	Dorsal tip of connecting process not pointed (figs. 13B,C); forearm 38-57 mm
4.	Antitragus nearly triangular (as in fig. 12A); cup at base of sella (immediately above nostrils) 8–10
	mm wide, almost as wide as anterior noseleaf; forearm 51-57 mm; skull as in Figure 45
	Rhinolophus philippinensis
4'.	Antitragus most nearly rectangular; cup at base of sella 2-4 mm wide, no more than half as wide
	as anterior noseleaf; forearm 38–56 mm
5.	Connecting process attaches below tip of sella (fig. 13B)
5'.	Connecting process attaches at tip of sella (fig. 13C)
6.	Ears 25-26 mm; sella 3-4 mm wide; forearm 43-44 mm; skull as in Figure 44 (see Notes)
6'.	Ears 17-21 mm; sella 1-2 mm wide; forearm 38-44 mm; skull as in Figure 46
7.	Forearm 43-46 mm; condylocanine length 16.2-17.4 mm; maxillary toothrow 6.7-7.5 mm; skull
	as in Figure 42 (see Notes)

TABLE 3. Measurement ranges of adult Philippine Rhinolophidae. Ranges represent at least 10 individuals, except when sample sizes are given in parentheses. Measurements, as defined in text, were taken from Philippine specimens unless otherwise indicated.

Species	CBL	CCL	C'-last M	Total Length	Tail	Hind Foot	Ear	Forearm	Weight
Coelops hirsuta*	(1) 13.0	ı	(1) 4.8	(1) 40	(1) 7	(1) 65	ı	(1) 34	1
Hipposideros ater	13.9-14.9	12.9-14.1	4.5-5.5	71-81	27–36	7-10	17-20	38-43	5-7
Hipposideros bicolor	(1) 16.2	(1) 15.2		ı	(1) 31	(1) 8	(1) 20	(1) 42	ı
Hipposideros bicolort	(5) 16.1–17.7	ı	(9) 5.6–6.2	74-85	25-32	8-9	18-20	41-47	(9) 7–10
Hipposideros cervinus	(1) 15.5	(1) 14.9		ı	ı	ı	1	ı	1
Hipposideros cervinus†‡	(7) 15.6–17.3	ı	5.8-6.5	77-81	(7) 24-32	(7) 8–9	(7) 11–16	44-50	(1) 7.5
Hipposideros coronatus*	ı	ı	(1) 7.0	(1) 100	(1) 34	(1) 10	(1) 15	(1) 47	ì
Hipposideros diadema (males)	26.1–29.9	24.9-28.2	10.5-12.0	140-151	44-50	16-21	30-35	79–89	34-84
Hipposideros diadema (females)	26.5-28.2	24.7–26.6	9.5-10.9	135-147	39-51	17-21	28-32	77–86	33–66
Hipposideros obscurus	15.4-17.2	14.7–16.7	5.8-6.7	71–79	18-24	10-12	18-22	42-48	7-12
Hipposideros pygmaeus	12.0-13.3	11.1–12.5	4.3-4.8	60–73	21–27	8-9	11-15	37-40	(3) 3-4
Rhinolophus acuminatus	19.5-20.5	(1) 18.3	7.4-8.1	(1) 85	(1) 25	(5) 11–13	(4) 16-20	(5)47-49	(3) 10-14
Rhinolophus arcuatus-s§	17.4-19.0	16.2-17.4	6.7-7.5	68-73	16-23	9-12	19-22	43-45.5	7-9.5
Rhinolophus arcuatus-1§	19.3-20.9	17.9-19.0	7.7-8.1	73-81	18-26	11-12	22-25	47-50	8.5-10
Rhinolophus inops	21.6-22.9	19.9-21.0	8.3-9.2	78–90	20-27	12-15	22-28	49-55	11-16
Rhinolophus macrotis	(3) 17.5 - 18.6	(4) 15.6–17.5	(8) 5.8–6.8	(2) 71–72	(2) 23-25	(2) 8-9	(2) 25-26	(2) 43-44	(1) 6
Rhinolophus philippinensis	(3) 21.7-22.2	(3) 19.9-20.3	(4) 7.9–8.2	(6) 89–94	(6) 29-33	(6) 10-12	(6) 28-36	(6) 51-57	(2) 11
Rhinolophus rufus	(6) 27.3–29.9	(6) 25.3–26.4	(6) 11.0–12.2	(8) 110-130	(7) 30-33	(7) 17-21	(8) 33–37	(8) 68-73	(8) 26–34
Rhinolophus subrufus	(6) 22.3–24.0	(1) 21.8	9.0-9.5	(8) 80–90	(9) 20-23	12-15	24-27	(9) 51–56	ı
Rhinolophus virgo	15.5-16.7	14.2-15.2	5.7-6.4	62–73	18-26	7-10	17-21	38-44	(6) 5–7

\* Measurements from Taylor, 1934.

† Borneo. ‡ Sulawesi. § See Notes section for explanation.

7'. Forearm 47-50 mm; condylocanine length 17.9-19.0 mm; maxillary toothrow 7.7-8.1 mm; skull 7". Forearm 49-55 mm; condylocanine length 19.9-21.0 mm; maxillary toothrow 8.3-9.2 mm; skull 7". Forearm 51-56 mm; condylocanine length about 21.8 mm; maxillary toothrow 9.0-9.5 mm (see Notes) Rhinolophus subrufus Tail absent; anterior noseleaf bilobed with two lateral leaflets underneath that project downward; no ridges on ears; well-developed secondary cusps on upper canines; forearm about 34 mm .... 8'. Tail present, greater than 15 mm; anterior noseleaf undivided, without leaflets that project downward; ridges on ears; only one cusp on upper canines; forearm 37-89 mm .... (Hipposideros) 9 9. Forearm 77-89 mm; cream-colored patch of fur anterior to leading edge of wing; skull as in Figure 11. Inner pair of lateral leaflets meet under anterior noseleaf but outer pair do not; forearm 37-40 mm; 11'. Both pairs of lateral leaflets do not meet under anterior noseleaf; forearm 44-50 mm ........ 12. No vertical septa in posterior noseleaf and tail > 30 mm; forearm about 47 mm (see Notes) .... 12'. Vertical septa may or may not be present in posterior noseleaf; if they are absent, then tail 18-24 13. Anterior noseleaf 5.5-7.0 mm wide, intermediate noseleaf 5.0-6.0 mm wide, posterior noseleaf 6.0-8.0 mm wide; hind foot length 10-12 mm; forearm 42-48 mm; skull as in Figure 39 ...... 13'. Anterior noseleaf 4.0-5.0 mm wide, intermediate noseleaf 3.5-4.0 mm wide, posterior noseleaf 14. Internarial septum not swollen at base; forearm about 42 mm .......... Hipposideros bicolor 14'. Internarial septum swollen at base; forearm 38-43 mm; skull as in Figure 37 ... Hipposideros ater

# Family Vespertilionidae: Evening Bats

This is an exceptionally large and diverse family, with about 40 genera and 275 species worldwide and about 11 genera and 22 species in the Philippines. Vespertilionids are identified as having a long tail that is completely enclosed by the interfemoral membrane (except occasionally the ex-




Fig. 13. Lateral views of connecting processes of some rhinolophids. A, R. acuminatus; B, R. macrotis (that of R. virgo is very similar); C, R. subrufus/R. arcuatus.

treme tip; fig. 7C), a simple face without a noseleaf (among species in the Philippines; fig. 8F), and a well-developed tragus. Most vespertilionids are small to medium-sized (forearm 22–55 mm; table 4) and have small eyes.

The skulls of vespertilionids are generally small, with a great variety of shapes and degrees of robustness (figs. 47–59). The braincase is high and domed in some genera (e.g., *Miniopterus, Myotis*; figs. 48–50, 52–55) and exceptionally low and flattened in others (e.g., *Tylonycteris*). They all lack postorbital processes. All species have a deep emargination at the anterior end of the palate, so that the incisors on either premaxillary are widely separated. The molariform teeth are generally typical of those of insectivorous bats, with broad crowns and sharp, shearing crests.

All of the vespertilionids in the Philippines are insectivorous. Roosting sites include caves, buildings, foliage, hollow trees, unfurled banana leaves,

TABLE 4. Measurement ranges of adult Philippine Vespertilionidae. Ranges represent at least 10 individuals, except when sample sizes are given in parentheses. Measurements, as defined in text, were taken from Philippine specimens unless otherwise indicated.

Species	CBL	CCL	C'-last M	Total Length	Tail	Hind Foot	Ear	Forearm	Weight
Glischropus tylopus*	(1) 11.4	10.7–11.6	4.3-4.5	70-90	33-40	9	10-11	26–30	1
Harpiocephalus harpia	(1) 20.6	(1) 19.7	(1) 6.9	(2) 121-125	(2) 46-51	(2) 13-14	(2) 18-19	(2) 51	19
Kerivoula hardwickii	(3) 12.6–12.7	(3) 11.9–12.4	(3) 5.1-5.4	(3) 77–88	(3) 38-47	(3) 8-10	(3) 14-15	(3) 34-35	(2) 4–5
Kerivoula pellucida	(1) 14.1	(1) 13.5	(1) 6.5	(1) 90	(1) 40	(1) 9	(1) 15	(1) 35	1
Kerivoula whiteheadi	(2) 11.8–12.1	(2) 11.3–11.4	(2) 5.2	(2) 66–72	(2) 31-37	(2) 7–8	(2) 12-14	(2) 30	1
Miniopterus australis	12.5-13.4	11.7–12.5	5.0-5.4	81-91	34-43	7–9	9-11	34-39	5-7
Miniopterus schreibersi	14.2-15.4	13.1-14.2	5.7-6.3	101-112	46-57	8-10	11-13	42-46	8-14
Miniopterus tristis	18.2-19.0	16.7-17.5	7.5-8.1	115-134	50-61	10-12	15-17	51-55	16-22
Murina cyclotis	(4) 16.6–17.5	(4) 15.8–16.8	(4) 5.9-6.3	(4) 90–101	(4) 38-42	(4) 9–12	(4) 14-17	(4) 36-39	(4) 9–11
Myotis horsfieldii	13.5-14.2	12.6-13.2	5.3-5.7	83-95	3041	10-12	12-16	35–38	5-6.5
Myotis macrotarsus	16.3-17.4	14.9–16.0	6.7 - 7.1	98-114	39-49	15-17	16-19	44-49	10-15
Myotis muricola	11.8-12.6	10.9-11.7	4.5-5.0	62-69	30–38	8-9	11-14	30–34	34
Myotis rufopictus	(4) 17.1 - 19.2	(4) 16.1–17.9	(4) 7.1 - 8.5	(1) 115	(1) 48	(1) 14	(1) 20	(2) 52–53	(1) 17
Philetor brachypterus	(7) 14.0–14.6	(7) 12.5–13.7	(7) 4.4-5.0	(7) 90–99	(7)31-39	(7) 9–11	(7) 13–15	(7)34-39	(7) 11-13
Phoniscus jagorii*+	(1) 15.4	ı	(2) 6.6-7.2	(2) 89–94	(2) 40-42	(2) 8-9	(1) 17	(1) 39	ı
Pipistrellus javanicus	12.6-13.6	11.8-12.8	4.5-5.0	75–90	29–37	7-10	10-13	32-36	4
Pipistrellus petersi	(2) 14.6 - 14.8	(2) 14.0	(2) 5.7	ı	ı	ı	1	(1) 38	ı
Pipistrellus stenopterus*;	(4) 15.1–15.9	(6) 14.9–15.3	(6) 5.7-6.1	(1) 92	(1) 36	(1) 11	(1) 13	(4)38-40	12-16
Pipistrellus tenuis	(6) 11.2–11.7	(6) 10.6–11.0	(6) 3.9-4.2	92–89 (9)	(6) 26-31	(4) 6–7	(6) 11-12	(4) 30-32	34
Scotophilus kuhlii	17.4-18.7	16.4–17.8	6.1–6.9	110-131	40-52	10-13	12-16	47-53	17–22
Tylonycteris pachypus	9.7-10.3	9.1–9.7	3.1–3.5	(2) 60–62	(2) 24-25	ı	(2) 7–8	22–25	1-2
Tylonycteris robustula	(4) 11.5-11.9	(4) 11.0-11.3	(4) 4.0-4.1	(2) 71–75	(2) 27	ı	(2) 8–9	(3) 24-27	I

\* Borneo. † Bali.

‡ Malaya.

and bamboo internodes. Some species have been captured only within forest, whereas others are common in urban areas. Species that roost in caves and buildings often form large colonies, ranging from hundreds to tens of thousands.

Many vespertilionid species that occur in the Philippines are described in Hill (1983). Other useful descriptions can be found in Francis and Hill (1986), Heaney and Alcala (1986), Hill (1963b, 1965, 1966, 1971), Kock (1981), and Tate (1942).

### Key to Vespertilionidae

1.	Nostrils elongated into short tubes that open laterally (fig. 14A); long orange hairs on dorsal surface of interfemoral membrane and feet; tragus straight and pointed, at least half length of ear (fig. 15A)
1′.	Nostrils not elongated into tubes (fig. 14B); hairs on interfemoral membrane very short and almost invisible; tragus shape and size varies between species
2.	Forearm 36–39 mm; five well-developed pairs of upper cheek teeth; skull as in Figure 51
2′.	Forearm about 51 mm; fifth pair of upper cheek teeth minute or absent (see Notes)
3.	Posterior margin of ear has a pronounced concave inflection near tip and forms a large, rounded flap below inflection, producing funnel-shaped ears (fig. 15B); tragus straight and pointed, at least half length of ear
3′.	No large, rounded flap on posterior margin of ear (figs. 15C,D); tragus shape and size varies between
4.	species
4'.	or salt-and-pepper appearance; forearm about 39 mm
5.	Bases of dorsal hairs pale; forearm about 35 mm
	Bases of dorsal hairs dark; forearm 30–35 mm
6.	Anterior two pairs of upper and lower premolars narrow and elongate, oval in cross-section; forearm
6′.	about 30 mm
	34–35 mm; skull as in Figure 47
7.	
	mm; skull as in Figure 59
	Two pairs of upper incisors; color of dorsal pelage varies between species; forearm 22–55 mm 8
8.	Tragus erect and tapering (fig. 15C)
	Tragus blunt (e.g., fig. 15D)
Э.	Wing membrane black except for reddish orange skin along digits; dorsal pelage reddish orange; forearm 52–53 mm; skull as in Figure 55
9'.	Wings uniformly colored; dorsal pelage gray or brown
	Posterior margin of wing membrane terminates on ankle (fig. 16A); hind foot 15-17 mm; forearm
	44–49 mm; skull as in Figure 53
10'.	Posterior margin of wing membrane terminates on side of foot, below ankle (figs. 16B,C); hind foot
	6–12 mm; forearm 30–38 mm
11.	Posterior margin of wing membrane terminates on side of foot, at least 1 mm above base of outermost toe (fig. 16B); hind foot 10-12 mm; forearm 35-38 mm; skull as in Figure 52
11'.	Posterior margin of wing membrane terminates on side of foot at base of outermost toe (fig. 16C);
	hind foot 6-8 mm; forearm 30-34 mm; skull as in Figure 54
12.	Pads present on feet and wrists (fig. 17); forearm 22–30 mm
	No pads on feet and wrists; forearm 30–55 mm

13. Pads on feet and wrists pink or white; dorsal fur dark at base; posterior margin of wing membrane




Fig. 14. Ventral views of muzzles of Murina (A) and Myotis (B).

terminates on side of foot, at base of outermost toe; five pairs of upper cheek teeth; forearm 26-30 mm Glischropus tylopus 13'. Pads on feet and wrists dark brown; dorsal fur not dark at base; posterior margin of wing membrane terminates on ankle or on side of foot, at least 1 mm above base of outermost toe; four pairs of upper cheek teeth; forearm 22–27 mm ...... (Tylonycteris) 14 14. Lower third of posterior margin of ear distinctly thicker than rest of ear (fig. 18A); forearm 24-27 14'. Lower third of posterior margin of ear not distinctly thicker than rest of ear (fig. 18B); forearm 22-15. Most distal bone of third digit of wing about three times length of adjoining phalanx ...... 15'. Most distal bone of third digit of wing (fig. 2) not more than two times length of adjoining phalanx 16. Forearm 51-55 mm; tail 50-61 mm; skull as in Figure 50 ...... Miniopterus tristis 17. Posterior margin of wing membrane terminates on lower leg, at or above ankle; fifth digit of wing (see fig. 2) does not extend beyond midpoint between first and second joint of third digit .... 18 17'. Posterior margin of wing membrane terminates on side of foot, below ankle (as in figs. 16B,C); fifth digit of wing extends beyond midpoint between first and second joint of third digit ........... 19 18. Inner upper incisors transversely elongate in cross-section, about three times as wide as outer upper incisors; outer secondary cusp on inner upper incisors is not as high but almost as wide as the main 18'. Inner upper incisors conical in shape, about two times as wide as outer upper incisors; outer




Fig. 15. Left external ears (pinnae) of some vespertilionids. A, Murina/Harpiocephalus; B, Kerivoula/Phoniscus; C, Myotis; D, Pipistrellus.




Fig. 16. Left hind feet of *Myotis* spp., showing attachment of wing membrane. A, *M. macrotarsus*; B, *M. horsfieldii*; C, *M. muricola*.

	secondary cusp on inner upper incisors is much narrower than main cusp; forearm 38–40 mm
	Pipistrellus stenopterus
19.	Forearm about 39 mm
19'.	Forearm 32–36 mm; tail 29–34 mm; skull as in Figure 57 Pipistrellus javanicus
19"	Forearm 30–32 mm: tail 26–31 mm: skull as in Figure 58 Pipistrellus tenuis

# Family Molossidae: Free-Tailed Bats

This family of at least 80 species is represented in the Philippines by only four species. As the English name for the family implies, molossid bats are the only bats with long tails that project well beyond the posterior edge of the interfemoral membrane (fig. 7E). The wings tend to be unusually long and narrow for the size of the bats. There is no noseleaf; the lips are often wrinkled and thick and the upper lip thickly sprinkled with short, bristly hairs (fig. 8E). The eyes are moderate to small. The external ear is moderately small, with the tragus small and the antitragus well developed. Philippine molossids are of moderate to large size, with forearms ranging from 40 to 90 mm (table

5). Two of the species in the Philippines (both *Cheiromeles*) have only scattered hairs over most of their bodies, leading to their common name, "naked bats."

The skulls of molossids are characterized by moderately sturdy construction, lack of postorbital processes, and broad, strong molars (fig. 60). Among Philippine molossids, the anterior edge of the palate is continuous, so that the upper pairs of incisors are separated by no more than a narrow gap. The jaws are powerful, and the large masseteric muscles on the head attach to a moderate to high sagittal crest.

These bats are strong, fast fliers that feed primarily on beetles and other large insects. *Chaere-phon plicata*, which roosts in caves and occasion-




Fig. 17. Pad on right wrist of Glischropus.




Fig. 18. Left ears of Tylonycteris. A, T. robustula; B, T. pachypus.

ally in buildings, probably once formed the largest colonies of bats in the Philippines, exceeding 100,000 individuals, according to early descriptions, but most colonies are now much reduced in size or entirely destroyed. *Cheiromeles parvidens* and *C. torquatus* are known to roost in hollow

coconut palms, hollow trees, and occasionally in caves, and *Mops sarasinorum* in hollow trees, where they form colonies of four to hundreds of individuals.

The systematics of bats of this family has been reviewed by Freeman (1981).

#### Key to Molossidae

- 3. Forearm 73–78 mm (occurs throughout the Philippines except Palawan) .... Cheiromeles parvidens

### Notes

The following brief comments on the status of several species of Philippine bats are intended to make users of the key aware of potential taxonomic problems and of recent records of bats that are referred to in the text.

### Acerodon lucifer

This species is known only from the type series collected on Panay Island in 1888. Based on current knowledge, it differs from most *Acerodon jubatus* only on the basis of smaller size, but it overlaps with some populations for all measurements. It is possible that careful study will show it to have defining characters that are not yet recognized, or, alternatively, to be a geographic variant of the widespread *A. jubatus*.

#### Cynopterus brachyotis

Philippine representatives of the genus *Cynopterus* have until recently been referred to *C. brachyotis*, which is widespread in Southeast Asia. Based primarily on morphometric analyses, however, Kitchener and Maharadatunkamsi (1990) split what had been known as *Cynopterus brachy*-

otis into several species, allocating Philippine specimens to *C. luzoniensis* based on a single sample from Negros Island. We have not yet been able to examine enough material to evaluate their proposed changes and have therefore followed the older terminology, while recognizing that further study may support the proposed revision.

#### Pteropus dasymallus

This species has been reported previously only from Japan, Taiwan, and adjacent and intervening islands. We have recently identified several specimens in FMNH, PNM, and USNM from the Babuyan and Batanes islands, which lie between northern Luzon and Taiwan, as representing this species. Details will be published elsewhere. The species is very similar to *P. hypomelanus*, differing most visibly in that the pelage is longer and denser, and the dorsal surface of the hind leg is thickly furred nearly to the ankle (rather than being nearly naked).

#### Hipposideros coronatus

No specimens of this moderately large horseshoe bat have been reported since the species was described in 1871 (Peters, 1871). Our placement

Measurement ranges of adult Philippine Molossidae. Ranges represent at least 10 individuals, except when sample sizes are given in parentheses. Measurements, as defined in text, were taken from Philippine specimens unless otherwise indicated. TABLE 5.

Species	CBL	CCL	C'-last M	Total Length	Tail	Hind Foot	Ear	Forearm	Weight
Chaerephon plicata	15.7–16.3	14.9–15.5	6.1–6.5	95–101	28–34	10-12	16-20	40-43	ı
Cheiromeles parvidens*	(1) 29.1	(2) 27.5-28.2	(2) 10.5–10.6	(4) 170 - 185	(4) 54-58	1	(4) 19-20	(4) 73-78	I
Cheiromeles torquatus†‡	(1) 29.9	(1) 28.4	(1) 11.5	(1) 261	(1) 59	(1) 26	(1) 29	80-90	1
Mops sarasinorum*	(7) 18.2–18.7	(7) 17.3–17.8	(7) 7.1–7.3	90-109	30–36	11-13	18-22	39-45	ı

External measurements from Taylor, 1934. External measurements taken from Bornean specimens.

External measurements taken from Bornean specimens. Forearm (of Thai bats) from Lekagul and McNeely, 1977.

of the species in the key is based solely on the original description and the comments of Taylor (1934).

### Rhinolophus anderseni

This species is known only from the holotype, which we have been unable to examine. As noted under the following species, it is possible that recently taken specimens may be referred to this species, but this remains uncertain.

### Rhinolophus arcuatus

Heaney et al. (1991) recently noted that bats previously identified as belonging to this species appear to fall into two morphotypes: one is slightly smaller with a narrower anterior noseleaf and is typically found in lowland caves or disturbed habitats, and the other is slightly larger with a proportionately wider anterior noseleaf and is usually found in primary upland forest. We designate these in the key as "R. arcuatus-s" and "R. arcuatus-l", respectively. J. Hill (in litt.) has suggested that one of these may be R. anderseni, but we have not examined the relevant holotypes and so remain uncertain of the appropriate use of names. More specimens, especially of the larger upland form, are badly needed.

# Rhinolophus inops and R. subrufus

Our studies of specimens currently in collections have indicated that Philippine bats in this species group may be broken into at least two morphs, based on size; these are very similar in morphology of the noseleaf and skull. In this key, we refer the range of smaller bats to *R. inops* and that of the larger ones to *R. subrufus*. However, we note that there are several unresolved problems with these bats.

First, the noseleaf of the holotype of *R. inops* has a distinctly shaped sella: the tip is modified into a downward-projecting triangular pouch (Andersen, 1905c). This modified sella is not found in any other specimen that has been referred to *R. inops* or in any other species in the genus; it may represent an aberration found in only one individual or it may be a diagnostic character for a species still known only from the holotype, with the other specimens referred to *R. inops* actually representing an undescribed species.

Second, the definition of *R. inops* is complicated by the variation in size among specimens that we have assigned to *R. inops*; available specimens from Negros are consistently smaller than those from Leyte and Biliran, whereas those from Catanduanes are intermediate (specimens from elsewhere are inadequate for comparison). It is possible that each morph represents a distinct species. Additional specimens and further study will be needed to clarify the status of all of these bats.

### Rhinolophus macrotis

The Philippine form of R. macrotis was first described as a distinct species, R. hirsutus (Andersen, 1905b), but was later subsumed under R. macrotis (Tate, 1943). We have examined the holotypes and referred specimens of both taxa and find that they differ in overall size, in proportionate tail length, and in the size and shape of noseleaf structures, particularly the sella. We suspect that the Philippine population is morphologically distinct and genetically independent and will eventually be shown to be a distinct species, but we refrain from making this change because we have not conducted comprehensive studies.

### Harpiocephalus harpia

Three specimens of this large and striking vespertilionid have been taken recently, one on Leyte (Rickart et al., in prep.), one on Luzon (Heaney et al., in prep.), and one on Negros (Utzurrum, pers. comm.). Further details will be published elsewhere.

### Acknowledgments

We wish to thank the curators and staff of the American Museum of Natural History, British Museum (Natural History), Delaware Museum of Natural History, Philippine National Museum, Royal Ontario Museum, Silliman University Museum of Natural History, University of Michigan Museum of Zoology, and United States National Museum of Natural History for allowing us to examine specimens in their care. We are particularly grateful to G. B. Corbet, P. C. Gonzales, P. Jenkins, K. F. Koopman, and W. T. Stanley for assistance with access to specimens. We have re-

ceived an exceptional amount of enthusiastic assistance with collecting bats in the field; although we are unable to mention all of those who have generously given us help, we must acknowledge the efforts of D. Balete, R. Fernandez, P. C. Gonzales, P. D. Heideman, A. Manamtam, E. A. Rickart, and R. C. B. Utzurrum. We are grateful to the Protected Areas and Wildlife Bureau of the Philippines, especially J. Caleda, C. Custodio, W. Dee, M. Mendoza, and S. Peñafiel for their continuing support and cooperation. The illustrations of skulls and the map were prepared by T. B. Griswold; all others were prepared by J. Sedlock, whom we especially thank for her efforts as a volunteer. C. M. Francis helped NRI distinguish between Philippine bats that also occur in Borneo. D. Balete, S. M. Goodman, K. F. Koopman, H. Miranda, A. T. Peterson, E. A. Rickart, W. Schutt, D. Willard, and an anonymous reviewer kindly reviewed the manuscript and provided helpful comments. We especially thank K. F. Koopman for his highly constructive input into every stage of this study. This research has been generously supported by the U.S. National Science Foundation (BSR-8514223) and the MacArthur Foundation (90-9272). The Dee Fund of the FMNH provided travel funds to NRI. An earlier draft of this manuscript formed part of a master's thesis by NRI at Cornell University.

#### Literature Cited

Andersen, K. 1905a. On some bats of the genus *Rhinolophus*, with remarks on their mutual affinities, and descriptions of twenty-six new forms. Proceedings of the Zoological Society of London, 2: 75–145.

——. 1905b. On the bats of the Rhinolophus philippinensis group, with descriptions of five new species. Annals and Magazine of Natural History, 7th series, 16: 243–256.

group, with descriptions of five new forms. Annals and Magazine of Natural History, 7th series, 16: 281-288.

tis group, with descriptions of two new forms. Annals and Magazine of Natural History, 7th series, 16: 289-292.

——. 1912. Catalog of the Chiroptera in the collection of the British Museum, vol. 1: Megachiroptera, 2nd ed. British Museum (Natural History), London, pp. ci + 854.

Anthony, E. L. P. 1988. Age determination in bats, pp. 47-58. *In* Kunz, T. K., ed., Ecological and Behavioral Methods for the Study of Bats. Smithsonian Institution Press, Washington, D.C.

Bergmans, W. 1975. A new species of *Dobsonia* Palmer, 1898 (Mammalia, Megachiroptera) from Waigeo,

- with notes on other members of the genus. Beaufortia, 23: 1–13.
- ——. 1978. On *Dobsonia* Palmer 1898 from the Lesser Sunda Islands (Mammalia: Megachiroptera). Senckenbergiana Biologica, **59**: 1–18.
- Collins, M., Ed. 1990. The Last Rainforests: A World Conservation Atlas. Oxford University Press, New York, 200 pp.
- DEBLASE, A. F., AND R. E. MARTIN. 1974. A Manual of Mammalogy with Keys to Families of the World. Wm. C. Brown Company Publishers, Dubuque, Iowa.
- Francis, C. M. 1989. Notes on fruit bats (Chiroptera, Pteropodidae) from Malaysia and Brunei, with the description of a new subspecies of *Megaerops wetmorei* Taylor, 1934. Canadian Journal of Zoology, **67**: 2878–2882.
- Francis, C. M., and J. E. Hill. 1986. A review of the Bornean *Pipistrellus* (Mammalia: Chiroptera). Mammalia, **50**: 43–55.
- Freeman, P. W. 1981. A multivariate study of the family Molossidae (Mammalia, Chiroptera): Morphology, ecology, evolution. Fieldiana, Zoology, n.s., 7: 1–173.
- HARRISON, J. 1966. An Introduction to the Mammals of Singapore and Malaysia. Malayan Nature Society, Singapore, 340 pp.
- HAUGE, P., J. TERBORGH, B. WINTER, AND J. PARKINSON. 1986. Conservation priorities in the Philippine Archipelago. Forktail, 2: 83–91.
- Heaney, L. R. 1986. Biogeography of mammals in Southeast Asia: Estimates of rates of colonization, extinction, and speciation. Biological Journal of the Linnean Society, 28: 127–165.
- ----. 1991. An analysis of patterns of distribution and species richness among Philippine fruit bats (Pteropodidae). Bulletin of the American Museum of Natural History, 206: 145–167.
- HEANEY, L. R., AND A. C. ALCALA. 1986. Flat-headed bats (Mammalia, *Tylonycteris*) from the Philippine Islands. Silliman Journal, **33**: 117–123.
- HEANEY, L. R., P. C. GONZALES, AND A. C. ALCALA. 1987. An annotated checklist of the taxonomic and conservation status of land mammals in the Philippines. Silliman Journal, 34: 32–66.
- HEANEY, L. R., P. C. GONZALES, R. C. B. UTZURRUM, AND E. A. RICKART. 1991. The mammals of Catanduanes Island: Implications for the biogeography of small land-bridge islands in the Philippines. Proceedings of the Biological Society of Washington, 104: 399–415.
- HEANEY, L. R., AND P. D. HEIDEMAN. 1987. Philippine fruit bats, endangered and extinct. Bats, 5: 3–5.
- Heaney, L. R., P. D. Heideman, E. A. Rickart, R. C. Utzurrum, and J. S. H. Klompen. 1989. Elevational zonation of mammals in the central Philippines. Journal of Tropical Ecology, 5: 259–280.
- Heaney, L. R., and R. L. Peterson. 1984. A new species of tube-nosed fruit bat (*Nyctimene*) from Negros Island, Philippines (Mammalia: Pteropodidae). Occasional Papers of the Museum of Zoology, University of Michigan, 708: 1–16.
- HEANEY, L. R., AND E. A. RICKART. 1990. Correlations

- of clades and clines: Geographic, elevational, and phylogenetic distribution patterns among Philippine mammals, pp. 321–332. *In* Peters, G., and R. Hutterer, eds., Vertebrates in the Tropics. Museum Alexander Koenig, Bonn, 424 pp.
- HEANEY, L. R., AND R. C. B. UTZURRUM, 1991. A Review of the Conservation Status of Philippine Land Mammals. Association of Systematic Biologists of the Philippines Communications (Manila), 3: 1–13.
- Heideman, P. D., and L. R. Heaney. 1989. Population biology of fruit bats (Pteropodidae) in Philippine submontane rainforest. Journal of Zoology (London), 218: 565–586.
- HILL, J. E. 1963a. A revision of the genus *Hipposideros*. Bulletin of the British Museum (Natural History), Zoology, 11: 1–129.
- ——. 1963b. Notes on some tube-nosed bats, genus *Murina*, from southeastern Asia, with descriptions of a new species and a new subspecies. Federation Museums Journal, new series, 8: 48–59.
- ——. 1965. Asiatic bats of the genera *Kerivoula* and *Phoniscus* (Vespertilionidae), with a note on *Kerivoula* aerosa Tomes. Mammalia, 29: 524–556.
- -----. 1966. A review of the genus *Philetor* (Chiroptera, Vespertilionidae). Bulletin of the British Museum (Natural History), Zoology, 14: 371–387.
- Temminck, 1840 (Chiroptera: Vespertilio brachypterus Zoologische Mededelingen, Leiden, 45: 139–146.
- ——. 1983. Bats (Mammalia: Chiroptera) from Indo-Australia. Bulletin of the British Museum (Natural History), Zoology, 45: 103–208.
- JENKINS, M. D., ED. 1987. Madagascar, an Environmental Profile. International Union for the Conservation of Nature, Gland, Switzerland. 374 pp.
- JENKINS, P. D., AND J. E. HILL. 1981. The status of Hipposideros galeritus Cantor, 1846 and Hipposideros cervinus (Gould, 1854) (Chiroptera: Hipposideridae). Bulletin of the British Museum (Natural History), Zoology, 41: 279-294.
- KITCHENER, D. J., AND MAHARADATUNKAMSI. 1990. Description of a new species of *Cynopterus* (Chiroptera: Pteropodidae) from Nusa Tenggara, Indonesia. Records of the Western Australian Museum, 15: 307–363.
- KLINGENER, D., AND G. K. CREIGHTON. 1984. On small bats of the genus *Pteropus* from the Philippines. Proceedings of the Biological Society of Washington, 97: 395–403.
- KOCK, D. 1969a. *Dyacopterus spadiceus* (Thomas 1890) auf den Philippinen (Mammalia, Chiroptera). Senckenbergiana Biologica, 50: 1–7.
- -----. 1969b. Eine neue Gattung und Art Cynopteriner Flughund von Mindanao, Philippinen (Mammalia, Chiroptera). Senckenbergiana Biologica, 50: 319-327.
- —. 1969c. Eine bemerkenswerte neue Gattung und Art Flughunde von Luzon, Philippinen (Mammalia, Chiroptera). Senckenbergiana Biologica, 50: 329–338.
- ——. 1981. Philetor brachypterus auf New-Britannien und den Philippinen (Mammalia: Chiroptera: Vespertilionidae). Senckenbergiana Biologica, 61: 313– 319.

- KOOPMAN, K. F. 1989. Distributional patterns of Indo-Malayan bats (Mammalia: Chiroptera). American Museum Novitates, 2942: 1-19.
- LAWRENCE, B. 1939. Collections from the Philippine Islands. Mammals. Bulletin of the Museum of Comparative Zoology at Harvard College, 86: 28–73.
- LEKAGUL, B., AND J. A. McNeely. 1977. Mammals of Thailand. Association for the Conservation of Wildlife, Bangkok, 758 pp.
- MEDWAY, L. 1969. The Wild Mammals of Malaya and Offshore Islands Including Singapore. Oxford University Press, London, 127 pp.
- MILLER, G. S. 1907. The families and genera of bats. Bulletin of the United States National Museum, 57: 1-282 + 14 pls.
- Musser, G. G., and L. R. Heaney. 1992. Philippine rodents: Definitions of *Tarsomys* and *Limnomys* plus a preliminary assessment of phylogenetic patterns among native Philippine murines (Murinae, Muridae). Bulletin of the American Museum of Natural History, 211: 1–138.
- Musser, G. G., K. F. Koopman, and D. Califia. 1982. The Sulawesian *Pteropus arquatus* and *P. argentatus* are *Acerodon celebensis*; the Philippine *P. leucotis* is an *Acerodon*. Journal of Mammalogy, **63**: 319–328.
- MYERS, N. 1988. Environmental degradation and some economic consequences in the Philippines. Environmental Conservation, 15: 205-214.
- Payne, J., C. M. Francis, and K. Phillipps. 1985. A Field Guide to the Mammals of Borneo. Sabah Society, Kota Kinabalu, 332 pp.
- Peters, W. 1871. Über die Gattungen und Arten der Huflisennasen, Rhinolophi. Monatsbericht der König-

- lich Preussischen Akademie der Wissenschaften zu Berlin, pp. 301-332.
- Peterson, R. L., and M. B. Fenton. 1970. Variation in the bats of the genus *Harpyionycteris*, with the description of a new race. Life Sciences Occasional Papers, Royal Ontario Museum, 17: 1–15.
- ROOKMAAKER, L. C., AND W. BERGMANS. 1981. Taxonomy and geography of *Rousettus amplexicaudatus* (Geoffroy, 1810) with comparative notes on sympatric congeners (Mammalia, Megachiroptera). Beaufortia, 31: 1–29.
- TATE, G. H. H. 1942. Results of the Archbold Expeditions. No. 47. Review of the vespertilionine bats, with special attention to genera and species of the Archbold Collections. Bulletin of the American Museum of Natural History, 80: 221–297.
- —. 1943. Results of the Archbold Expeditions. No. 49. Further notes on the *Rhinolophus philippinensis* group (Chiroptera). American Museum Novitates, 1219: 1–5.
- TAYLOR, E. H. 1934. Philippine land mammals. Monographs of the Bureau of Science, Manila, 30: 1–548 + 25 pls.
- UTZURRUM, R. C. B. 1991. Philippine island biogeographic patterns: Practical applications for resource conservation and management. Association of Systematic Biologists of the Philippines Communications (Manila), 3: 19–32.
- YOSHIYUKI, M. 1979. A new species of the genus *Ptenochirus* (Chiroptera, Pteropodidae) from the Philippine Islands. Bulletin of the National Science Museum, Tokyo, series A (Zoology), 5: 75–81.




Fig. 19. Skull of Acerodon jubatus (USNM 74321). Scale = 10 mm.




Fig. 20. Skull of Cynopterus brachyotis (UMMZ 156664). Scale = 10 mm.




Fig. 21. Skull of Dobsonia chapmani (DMNH 5131). Scale = 10 mm.




Fig. 22. Skull of Eonycteris robusta (UMMZ 162221). Scale = 10 mm.




Fig. 23. Skull of *Eonycteris spelaea* (UMMZ 158482). Scale = 10 mm.




Fig. 24. Skull of *Haplonycteris fischeri* (UMMZ 157047). Scale = 10 mm.




Fig. 25. Skull of *Harpyionycteris whiteheadi* (UMMZ 158838). Scale = 10 mm.




Fig. 26. Skull of Macroglossus minimus (UMMZ 158725). Scale = 10 mm.




Fig. 27. Skull of Nyctimene rabori (USNM 458281). Scale = 10 mm.




Fig. 28. Skull of Ptenochirus jagori (UMMZ 158513). Scale = 10 mm.




Fig. 29. Skull of Pteropus hypomelanus (USNM 105443). Scale = 10 mm.




Fig. 30. Skull of *Pteropus pumilus* (UMMZ 158517). Scale = 10 mm.




Fig. 31. Skull of Pteropus vampyrus (UMMZ 158849). Scale = 10 mm.




Fig. 32. Skull of Rousettus amplexicaudatus (UMMZ 162304). Scale = 10 mm.




Fig. 33. Skull of *Emballonura alecto* (USNM 458536). Scale = 10 mm.




Fig. 34. Skull of Saccolaimus saccolaimus (USNM 458555). Scale = 10 mm.




Fig. 35. Skull of *Taphozous melanopogon* (USNM 458571). Scale = 10 mm.




Fig. 36. Skull of Megaderma spasma (UMMZ 160295). Scale = 10 mm.




Fig. 37. Skull of *Hipposideros ater* (ROM 40735). Scale = 10 mm.




Fig. 38. Skull of *Hipposideros diadema* (UMMZ 160296). Scale = 10 mm.




Fig. 39. Skull of *Hipposideros obscurus* (su 787). Scale = 10 mm.




Fig. 40. Skull of *Hipposideros pygmaeus* (UMMZ 156872). Scale = 10 mm.




Fig. 41. Skull of Rhinolophus arcuatus-l (UMMZ 158526). Scale = 10 mm.




Fig. 42. Skull of Rhinolophus arcuatus-s (UMMZ 157106). Scale = 10 mm.




Fig. 43. Skull of Rhinolophus inops (USNM 459495). Scale = 10 mm.




Fig. 44. Skull of Rhinolophus macrotis (UMMZ 160328). Scale = 10 mm.




Fig. 45. Skull of Rhinolophus philippinensis (UMMZ 459497). Scale = 10 mm.




Fig. 46. Skull of Rhinolophus virgo (USNM 459454). Scale = 10 mm.




Fig. 47. Skull of Kerivoula hardwickii (USNM 459511). Scale = 10 mm.




Fig. 48. Skull of Miniopterus australis (USNM 458658). Scale = 10 mm.




Fig. 49. Skull of Miniopterus schreibersi (UMMZ 157001). Scale = 10 mm.




Fig. 50. Skull of *Miniopterus tristis* (USNM 458680). Scale = 10 mm.




Fig. 51. Skull of Murina cyclotis (USNM 573776). Scale = 10 mm.




Fig. 52. Skull of Myotis horsfieldii (UMMZ 158872). Scale = 10 mm.




Fig. 53. Skull of Myotis macrotarsus (UMMZ 162364). Scale = 10 mm.




Fig. 54. Skull of *Myotis muricola* (UMMZ 158873). Scale = 10 mm.




Fig. 55. Skull of Myotis rufopictus (UMMZ 158878). Scale = 10 mm.




Fig. 56. Skull of *Philetor brachypterus* (USNM 573781). Scale = 10 mm.




Fig. 57. Skull of Pipistrellus javanicus (USNM 459762). Scale = 10 mm.




Fig. 58. Skull of *Pipistrellus tenuis* (USNM 160338). Scale = 10 mm.




Fig. 59. Skull of Scotophilus kuhlii (USNM 458704). Scale = 10 mm.




Fig. 60. Skull of Chaerephon plicata (USNM 304238). Scale = 10 mm.